

# Communicable Disease Reference Chart for School and Child Care Facility Personnel\*

DISEASE	INCUBATION PERIOD	TRANSMISSION	COMMON SYMPTOMS	RECOMMENDATIONS
<a href="#">Chickenpox</a> <sup>†</sup> (Varicella)	14–16 days (Range 10–21 days)  Incubation period may be longer (28 days or more) in persons who receive immune globulin for postexposure prophylaxis (VariZIG or IVIG). Incubation period may be shorter in persons with weakened immune systems.	Direct contact with vesicular fluid or by airborne spread from respiratory tract secretions.  Infectious from 1–2 days before rash onset until all lesions have dried/crusted over and no new lesions appear within a 24-hour period (average is 4–7 days). Communicability may be prolonged in people with weakened immune systems.	Sudden onset with slight fever, other systemic symptoms and itchy eruptions which become vesicular (small blisters) within a few hours. The rash first appears on the chest, back, and face, and then spreads over the entire body. Lesions commonly occur in successive crops, with several stages of maturity present at the same time. Typically, vesicular rash consisting of 250–500 lesions in varying stages of development (papules, vesicles) and resolution (crusting).	<b>PATIENT:</b> Exclude from school or child care until: <ul style="list-style-type: none"> <li>Uncomplicated varicella: the rash has crusted</li> <li>Immunized patients without crust: no new lesions appear within a 24-hour period</li> <li>Patients with weakened immune systems: at least 5 days after eruptions first appear or until vesicles become dry</li> </ul> Avoid exposure to women in early pregnancy who have not had chickenpox and/or varicella vaccine.  <b>CONTACTS:</b> Check vaccination status of contacts in school and child care and recommend vaccination if needed within 3 days and up to 5 days after exposure. For exposed contacts without immunity: <ul style="list-style-type: none"> <li><u>School</u> setting: exclusion not recommended unless an outbreak is reported; monitor for varicella symptoms for 8–21 days from exposure and until 28 days after for those who received VariZIG or IVIG.</li> <li><u>Child care</u> setting: exclusion recommended for 8–21 days from exposure (28 days if VariZIG or IVIG given.)</li> </ul> If symptoms occur, exclude from school and child care.
<a href="#">Conjunctivitis</a> (Pink Eye)	1–3 days Variable depending on the cause.  Causes include bacteria, viruses, allergies, and chemicals/irritants.	Contact with eye discharge or contaminated articles, if the cause is infectious (e.g., bacteria or virus.)  Allergic and chemical causes are not contagious.	Pink or red eye with swelling of the eyelids, increased tears, and eye discharge. Eyelids may be matted shut after sleep. May involve one or both eyes.	<b>PATIENT:</b> Exclude from school or child care while symptomatic or until cleared to return by a healthcare provider. Important to wash hands thoroughly after contact with eye drainage. Do not share any articles that have come into contact with the eyes.  <b>CONTACTS:</b> Exclusion not indicated.
<a href="#">COVID-19</a> <sup>†</sup> (Coronavirus disease 2019 caused by SARS-CoV-2 virus)	2–14 days  As virus variants continue to emerge, characteristics such as incubation period and ease of transmission may change.	Inhalation or contact with respiratory droplets; contact with contaminated surfaces.  People are infectious 2 days before symptom onset through 10 days after symptom onset, with viral loads higher earlier in the course of infection.	Patients may be asymptomatic or may have a wide variety of symptoms of variable severity: <ul style="list-style-type: none"> <li>Fever and/or chills</li> <li>Cough and/or runny nose</li> <li>Shortness of breath</li> <li>Headache</li> <li>Fatigue</li> <li>Sore throat</li> <li>Muscle or body aches</li> <li>New loss of taste or smell</li> <li>Nausea, vomiting, diarrhea</li> </ul>	<b>PATIENT:</b> Please refer to CDC’s Preventing Spread of Infections in K-12 Schools “ <a href="#">When Students or Staff are Sick</a> .” Depending on severity of illness, refer to a healthcare provider, urgent care center, or emergency department for further evaluation and treatment.  <b>CONTACTS:</b> Quarantine is not recommended for contacts.

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

\*Please see last page for a summary of major changes.

<sup>†</sup>Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>

# Communicable Disease Reference Chart for School and Child Care Facility Personnel\*

DISEASE	INCUBATION PERIOD	TRANSMISSION	COMMON SYMPTOMS	RECOMMENDATIONS
<p>Diarrheal Diseases† (<a href="#">Campylobacteriosis</a>, <a href="#">E. coli O157:H7</a>, <a href="#">Giardiasis</a>, <a href="#">Salmonellosis</a>, <a href="#">Shigellosis</a>, etc.)</p>	<p>Campylobacteriosis: 2–5 days (Range: 1–10 days)</p> <p><i>E. coli</i> O157:H7: 3–4 days (Range: 1–10 days)</p> <p>Giardiasis: 1–3 weeks</p> <p>Salmonellosis: 12–36 hours (Range: 6–72 hours; though up to 16 days has been reported)</p> <p>Shigellosis: 1–3 days (Range: 1–7 days)</p>	<p>Fecal-oral route through direct contact or by ingestion of contaminated, raw, or improperly cooked food or untreated water</p>	<p>The most common symptoms are diarrhea, abdominal pain or cramping, malaise, and fever. Stools can contain visible or occult blood. Severe dehydration may occur in young children.</p> <p>Campylobacteriosis may mimic appendicitis or intussusception. <i>E. coli</i> O157:H7 may lead to hemolytic-uremic syndrome. In giardiasis, persons may be asymptomatic or have malabsorption with smelly, greasy stool and weight loss.</p> <p>Patients with weakened immune systems can have prolonged, relapsing, or extraintestinal infections.</p>	<p><b>PATIENT:</b> Exclude from school and child care until stools are contained in the diaper or when continent patients no longer have fecal accidents and when stool frequency becomes no more than 2 stools above normal frequency for the patient, even if the stools remain loose. Stress importance of proper handwashing. In an outbreak setting, consult local health department for clearance to return to school/childcare setting. Please note that other resources advise no return to school until diarrhea has ceased for at least 24 hours.</p> <p><b>CONTACTS:</b> Exclusion and stool cultures not indicated in absence of symptoms. Contacts who are symptomatic should be excluded until stools are contained in the diaper or child is continent and stool frequency is no more than 2 stools above that child's normal frequency for the time the child is in the program. Stool cultures are recommended for symptomatic contacts, and these children should be excluded from school and child care while evaluation is pending.</p> <p>Consult with your local health department for advice during suspected school outbreaks.</p> <p>Thorough hand hygiene and environmental cleaning is very important.</p>
<p><a href="#">Fifth Disease</a> (Parvovirus B19, Erythema Infectiosum)</p>	<p>4–14 days (Range: 4–21 days)</p>	<p>Contact with respiratory tract secretions and percutaneous exposure to blood or blood products. It can also be transmitted from mother to baby during pregnancy.</p> <p>Once the characteristic rash appears, the child is not likely to be contagious and can return to school.</p>	<p>Distinctive rash characterized by a vivid reddening of the skin, especially of the face, which fades and recurs; classically, described as a “slapped cheek appearance.” The rash may have a lacey appearance on the trunk and extremities. It can fluctuate in intensity and recur with environmental changes, such as temperature and exposure to sunlight, for weeks to months. Mild symptoms of fever, joint pain, cough, sore throat, and headache may occur 7–10 days before rash.</p>	<p><b>PATIENT:</b> Exclusion not indicated beyond facility-based policies about fever.</p> <p><b>CONTACTS:</b> Exclusion not indicated. Pregnant women and persons with weakened immune systems should seek medical advice.</p>

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

\*Please see last page for a summary of major changes.

†Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>

# Communicable Disease Reference Chart for School and Child Care Facility Personnel\*

DISEASE	INCUBATION PERIOD	TRANSMISSION	COMMON SYMPTOMS	RECOMMENDATIONS
<p><a href="#">Hepatitis A</a><sup>†</sup></p> <p><b>All suspected or confirmed cases of hepatitis A are rapidly reportable to the local health department</b></p>	<p>28 days (Range: 15–50 days)</p>	<p>Fecal-oral route through direct contact or ingestion of contaminated food or water.</p> <p>People with Hepatitis A virus (HAV) infection are most infectious during the 1–2 weeks before illness onset, when concentration of virus in the stool is highest. Risk diminishes significantly by 7–10 days after symptom onset and is minimal by the 3<sup>rd</sup> week of illness.</p>	<p>Initial symptoms begin abruptly and include fever, nausea, vomiting, diarrhea, fatigue, joint pain, anorexia, and abdominal pain or discomfort. Dark urine, pale stools, and jaundice (yellowing of the skin or eyes) might be present initially or might develop a few days to a week later. The likelihood of symptoms increases with age.</p> <p>Symptoms typically last less than 2 months but can extend up to 6 months.</p>	<p><b>PATIENT:</b> Patients with acute HAV infection who work as food handlers or attend or work in child care settings should be excluded for 14 days if symptomatic, or 7 days since jaundice. If the person is not a food handler or in a child care setting, exclude for 7 days after onset of symptoms. Serologic testing should be performed to confirm HAV infection in suspected cases.</p> <p><b>CONTACTS:</b> Determine if contact is immune to HAV through immunization records and/or serologic testing (a positive Hep A IgG). Serologic testing is not typically recommended if immunization records are unavailable. Contacts who are immune do not need additional follow-up. Determine if contact meets criteria for HAV postexposure prophylaxis (PEP) with Hepatitis A vaccine, immunoglobulin, or both.</p> <p>Childcare staff or attendees who might have been exposed to HAV in the past 2 weeks should be excluded for 7 weeks after last contact with the case during the infectious period, unless they receive PEP within 14 days of exposure or report immunity through vaccination or serology indicating previous disease.</p> <p>Contacts who are symptomatic and epidemiologically-linked to a laboratory-confirmed case meet the case definition and should be managed as a case, including appropriate investigation and exclusion.</p> <p>In a child care center, stress importance of proper handwashing. In facilities with diapered children, if one or more cases are confirmed in child or staff attendees or two or more cases in households of staff or attendees, hepatitis A PEP should be administered within 14 days of exposure to all unimmunized staff and attendees.</p> <p>In centers without diapered children, hepatitis A PEP should be administered only to unimmunized classroom contacts of index case. Asymptomatic contacts may return after receipt of hepatitis A PEP.</p> <p>If questions about management of cases and/or contacts, please contact local health department.</p>

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

\*Please see last page for a summary of major changes.

<sup>†</sup>Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>

# Communicable Disease Reference Chart for School and Child Care Facility Personnel\*

DISEASE	INCUBATION PERIOD	TRANSMISSION	COMMON SYMPTOMS	RECOMMENDATIONS
<a href="#">Hepatitis B†</a>	60–90 days (Range: 45–160 days)	Direct contact with infected blood or body fluids. Transmission occurs when the hepatitis B virus enters the body through broken skin or mucous membranes. The risks of HBV acquisition when a susceptible child bites a child who has chronic HBV infection or when a susceptible child is bitten by a child with chronic HBV infection are unknown. A theoretical risk exists if HBsAg-positive blood enters the oral cavity of the biter, but transmission by this route has not been reported.	Developing symptoms of acute hepatitis are age-dependent and occur most commonly in adults. The spectrum of symptoms and signs is varied and includes subacute illness with nonspecific symptoms (e.g., anorexia, nausea, or malaise), clinical hepatitis with jaundice, or fulminant hepatitis.	PATIENT: Follow advice of child’s healthcare provider and/or your local health department.  CONTACTS: Exclusion not indicated.
<a href="#">HIV Infection† and AIDS†</a> (Acquired Immunodeficiency Syndrome)	For untreated infants and children who acquire HIV infection through mother-to-child transmission, the usual age of onset of symptoms is 12–18 months of age. Some children become ill in the first few months of life, but others remain asymptomatic for more than 5 years and, rarely, until early adolescence.  Acute retroviral syndrome develops in 40–90% of adults and adolescents. It occurs 7–14 days following viral acquisition and lasts 5–7 days. Most patients are not ill enough to seek medical attention.	<ol style="list-style-type: none"> <li>1. Sexual transmission (vaginal, anal, orogenital)</li> <li>2. Percutaneous blood exposure (e.g., contaminated needles)</li> <li>3. Mother-to-child transmission</li> <li>4. Mucous membrane exposure to contaminated blood or body fluids</li> <li>5. Transfusion of contaminated blood or blood products, though this has been virtually eliminated in the U.S. since 1985</li> <li>6. Cases of HIV transmission have been reported from contact of non-intact skin with blood-containing body fluids</li> </ol>	Clinical manifestations of untreated pediatric HIV infection include unexplained fevers, generalized lymphadenopathy, hepatomegaly, splenomegaly, failure to thrive, persistent oral and diaper candidiasis, recurrent diarrhea, parotitis, hepatitis, central nervous system disease (e.g., encephalopathy, hyperreflexia, hypertonia, floppiness, developmental delay), lymphoid interstitial pneumonia, recurrent invasive bacterial infections, and opportunistic infections (e.g., viral, parasitic, and fungal infections).	PATIENT: Follow advice of child’s healthcare provider and/or your local health department.  CONTACTS: Exclusion not indicated.

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

\*Please see last page for a summary of major changes.

†Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>

# Communicable Disease Reference Chart for School and Child Care Facility Personnel\*

DISEASE	INCUBATION PERIOD	TRANSMISSION	COMMON SYMPTOMS	RECOMMENDATIONS
<p><a href="#">Influenza</a> (Seasonal)</p> <p>Note: Suspected or confirmed novel influenza virus infections should be rapidly reported to the local health department and may have different exclusion criteria.</p>	<p>2 days (Range: 1–4 days)</p>	<p>Respiratory droplets (e.g., coughing or sneezing.), contaminated hands or surfaces (where it can remain for up to 24 hours), or airborne transmission via small-particle aerosols in the vicinity of the infectious individual</p> <p>Patients may be infectious 24 hours before onset of symptoms. Viral shedding in nasal secretions usually peaks during the first 3 days of illness and ceases within 7 days but can be prolonged (10 days or longer) in young children and people with weakened immune systems.</p>	<p>Sudden onset of fever, often accompanied by nonproductive cough, chills or rigors, diffuse myalgia, headache, and malaise. Subsequently, respiratory tract symptoms, including sore throat, nasal congestion, rhinitis, and cough, become more prominent. Less commonly, abdominal pain, nausea, vomiting, and diarrhea are associated with influenza illness. In some children, influenza can appear as an upper respiratory tract illness or as a febrile illness with few respiratory tract symptoms. In infants, influenza can produce a nonspecific sepsis-like illness picture, and in infants and young children, influenza can cause otitis media, croup, pertussis like-illness, bronchiolitis, or pneumonia. Acute myositis secondary to influenza can present with calf tenderness and refusal to walk.</p>	<p>PATIENT: Exclude from school and child care until at least 24 hours following resolution of fever without the use of fever-reducing medication(s).</p> <p>CONTACTS: Exclusion not indicated.</p> <p>Annual seasonal influenza vaccination for staff and children ≥ 6 months of age is strongly encouraged to prevent cases of influenza or lessen severity of illness.</p>
<p><a href="#">Measles</a><sup>†</sup> (Rubeola, Red Measles)</p> <p><b>All suspected or confirmed measles cases are rapidly reportable to the local health department</b></p>	<p>From exposure to prodrome: 11–12 days</p> <p>From exposure to rash onset: 14 days (Range: 7–21 days)</p>	<p>Direct contact with infectious droplets or by airborne spread through inhalation of infectious droplets when a person with measles coughs, sneezes, etc. The attack rate in a susceptible individual exposed to measles is 90% in close-contact settings.</p> <p>Patients infected with wild-type measles virus are contagious from 4 days before rash onset through 4 days after appearance of the rash.</p>	<p>Fever, cough, coryza, and conjunctivitis, followed by a maculopapular rash starting on the face and spreading downwards and outwards. During the initial period, Koplik spots may be present. The characteristic rash may not develop in patients with weakened immune systems.</p> <p>Koplik spots are typically 1-3mm whitish, grayish, or bluish elevations with an erythematous base typically seen on the buccal mucosa. Koplik spots are not seen in all measles cases.</p>	<p>PATIENT: Exclude from school and child care until at least 4 days after appearance of the rash and when the child is able to participate.</p> <p>CONTACTS: Check immunization records of all contacts. Additional measures are not needed for children and/or staff who are fully vaccinated or have evidence of immunity to measles. People who have not been immunized, including those exempted from measles immunization for medical reasons, should be excluded from school, child care, and health care settings until at least 21 days after the onset of rash in the last case of measles. Discuss with your local health department. Every suspected measles case should be reported immediately to the local health department.</p>

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

\*Please see last page for a summary of major changes.

†Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>

# Communicable Disease Reference Chart for School and Child Care Facility Personnel\*

DISEASE	INCUBATION PERIOD	TRANSMISSION	COMMON SYMPTOMS	RECOMMENDATIONS
<p><a href="#">Meningitis, Bacterial</a> (<a href="#">H. influenzae</a><sup>†</sup>, <a href="#">Meningococcal</a><sup>†</sup>, <a href="#">Pneumococcal</a>)</p> <p><b>All suspected or confirmed cases of invasive H. influenzae disease or meningococcal disease are rapidly reportable to the local health department</b></p>	<p><i>H. influenzae</i>: 2–7 days</p> <p>Meningococcal: &lt; 4 days (Range: 1–10 days)</p> <p>Pneumococcal: 1–7 days</p>	<p>Respiratory (e.g., coughing or sneezing) or throat secretions (e.g., kissing) through close or lengthy contact.</p> <p>Transmission can also occur by sharing beverages, utensils, and personal care products. Patients should be considered capable of transmitting the organism for up to 24 hours after initiation of effective antimicrobial treatment.</p>	<p>Acute onset of fever, neck stiffness, neck pain, headache, light sensitivity and other neurologic symptoms or signs.</p> <p>In meningococcal disease, rash initially can be macular or maculopapular but typically becomes petechial or purpuric within hours. Signs and symptoms of meningococcal meningitis may be indistinguishable from those associated with pneumococcal meningitis.</p>	<p><b>PATIENT:</b> Exclude from school and child care during acute illness. Non-communicable after 24–48 hours of appropriate drug therapy.</p> <p><b>CONTACTS:</b> Exclusion not indicated. Discuss with your local health department to determine if close contacts need <b>prompt prophylactic treatment</b> for <i>H. influenzae</i> or meningococcal meningitis, especially contacts in a child care setting. Students and staff should be advised not to share beverages, eating utensils, or personal care products.</p>
<p><a href="#">Mpox</a><sup>†</sup> (Previously known as monkeypox)</p>	<p>3–17 days</p> <p>Symptoms usually start within 21 days of exposure to the virus. If someone has flu-like symptoms, they will usually develop a rash 1–4 days later.</p>	<p>Through close contact (including intimate, sexual, or household/caregiving contact) with a person with mpox (e.g., skin-to-skin, body fluids, prolonged talking or breathing, from mother to baby), and through contact with contaminated materials (e.g., clothing, bedding), or direct contact with infected wild animals.</p> <p>Mpox can be spread from the time symptoms start until the rash has fully healed and a fresh layer of skin has formed. The illness typically lasts 2–4 weeks.</p>	<p>People with mpox get a rash that may be located on the hands, feet, chest, face, in or around the mouth, or on or around the genitals or anus.</p> <p>The rash will go through several stages, including scabs, before healing. The rash can initially look like pimples or blisters and may be painful or itchy.</p> <p>Other symptoms of mpox can include fever, chills, swollen lymph nodes, exhaustion, muscle aches and backache, headache, respiratory symptoms (e.g., sore throat, nasal congestion, or cough).</p> <p>People may have all or only a few symptoms.</p>	<p><b>PATIENT:</b> Exclude from school and child care until the rash has healed, all scabs have fallen off, and a fresh layer of skin has formed. This may take as long as 4 weeks after symptoms began. Caregivers should work with a healthcare provider and the health department to decide when the child or adolescent can return to the educational setting.</p> <p><b>CONTACTS:</b> Exclusion not indicated in most cases. In some cases, if contact tracing may not be possible and there was a high or intermediate degree of exposure, the local health department may consider limiting an individual’s participation in activities. The health department will consider the age of the individual and their ability to recognize or communicate symptoms, the types of interactions in the environment, and the risk of more severe disease to others in the setting. If a contact had high- or intermediate-risk exposure, the health department may recommend postexposure prophylaxis with JYNNEOS vaccine, ideally administered within 4 days of exposure.</p> <p>Settings with children or adolescents in residence, (e.g., boarding schools, overnight camps, and other residential environments) follow considerations for congregate settings.</p>

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

<sup>†</sup>Please see last page for a summary of major changes.

\*Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>

# Communicable Disease Reference Chart for School and Child Care Facility Personnel\*

DISEASE	INCUBATION PERIOD	TRANSMISSION	COMMON SYMPTOMS	RECOMMENDATIONS
<a href="#">Mumps</a> <sup>†</sup>	16–18 days (Range 12–25 days)	<p>Contact with infectious respiratory tract secretions and saliva. Highly contagious.</p> <p>The period of maximum communicability begins 2 days before parotitis onset. The recommended isolation period is 5 days after onset of parotid swelling. However, virus has been detected in patients' saliva as early as 7 days before and until 9 days after onset of swelling.</p>	Swelling and tenderness of one or both parotid glands as well as nonspecific symptoms such as fever, muscle aches, loss of appetite, malaise, and headache. Asymptomatic infection occurs in about 15-20% of cases, usually in adults more than children.	<p><b>PATIENT:</b> In addition to standard precautions, exclude from school and child care for 5 days after the onset of parotid gland swelling.</p> <p><b>CONTACTS:</b> Determination of vaccination status should be obtained for all contacts. In an outbreak setting, consult local health department regarding outbreak management and whether contacts need to be excluded from school or child care.</p> <p>If health department agrees, unimmunized contacts should be excluded until at least 26 days after onset of parotitis in the last person with mumps. Excluded students can be readmitted after receipt of a dose of MMR vaccine at the discretion of the health department. Moreover, if the school or child care is considered to be at high risk for mumps, a recommendation may be made for students and staff who have had two doses of a mumps-containing vaccine to receive a third dose of MMR vaccine.</p>
<a href="#">Norovirus</a>	12–48 hours	<p>Fecal-oral or vomitus-oral route through direct contact, ingestion of contaminated food or water, touching contaminated surfaces, and aerosolized vomitus.</p> <p>Viral shedding may start before onset of symptoms, peaks several days after exposure, and in some cases, may persist for ≥4 weeks. Prolonged shedding (&gt;6 months) has been reported in people with weakened immune systems.</p>	Sudden onset vomiting and/or watery diarrhea, abdominal cramps, and nausea. Symptoms typically last from 24–72 hours, but prolonged illness can occur. Systemic manifestations, including fever, myalgia, malaise, anorexia, and headache, may accompany gastrointestinal tract symptoms.	<p><b>PATIENT:</b> In addition to standard precautions, exclude from school and child care until 48 hours after symptoms resolve. This includes resolution of vomiting and diarrhea. Infants and children should be excluded from child care centers until stools are contained in the diaper or when toilet-trained children no longer have accidents using the toilet and when stool frequency becomes no more than 2 stools above that child's normal frequency for the time the child is in the program, even if the stools remain loose. Stress importance of thorough handwashing with soap and water, especially when caring for diapered children, as virus may be shed in stool for weeks after symptoms resolve.</p> <p><b>CONTACTS:</b> Exclusion not indicated.</p> <p>Environmental cleaning is a very important component of the response to a norovirus outbreak. A high-concentration bleach solution can be used—this solution must remain on the surface for enough time to kill norovirus. Surface must be rinsed with water to remove bleach residue.</p>

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

\*Please see last page for a summary of major changes.

<sup>†</sup>Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>

# Communicable Disease Reference Chart for School and Child Care Facility Personnel\*

DISEASE	INCUBATION PERIOD	TRANSMISSION	COMMON SYMPTOMS	RECOMMENDATIONS
<a href="#">Pediculosis</a> (Head Lice)	8–9 days for eggs to hatch after being laid (Range 7–12 days)  Lice mature into adults about 7 days later.	Direct head-to-head contact with hair of an infected person. Transmission by contact with personal belongings (e.g., combs, brushes, sporting gear, hats) is uncommon.  Head lice infestation occurs most commonly in children attending child care, preschool, and elementary school, and is not a sign of poor hygiene.	Severe itching and scratching. Adults, eggs, and nits (empty egg casings) are found on the hair, often seen behind ears and at the nape of the neck. Excoriations and crusting caused by secondary bacterial infection may occur.  Many children are asymptomatic.	<b>PATIENT:</b> Notify parents; inform that child has lice and should be treated. Children should not be excluded or sent home early from school or child care because of head lice, because head lice have a low contagion within classrooms. “No-nit” policies should be discouraged.  <b>CONTACTS:</b> Household and close contacts should be examined and treated if infested. No exclusion necessary. Routine classroom or school-wide screening for lice is discouraged. Parents who are educated on the diagnosis of lice infestation may screen their own children’s heads for lice regularly and if the child is symptomatic. Refer for treatment if infested.
<a href="#">Pertussis</a> <sup>†</sup> (Whooping Cough)  <b>All suspected or confirmed cases of pertussis are rapidly reportable to the local health department</b>	7–10 days (Range 5–21 days)	Close contact through large respiratory droplets generated by coughing or sneezing.	<b>Catarrhal stage:</b> mild upper respiratory tract symptoms (e.g., common cold). Lasts 1–2 weeks.  <b>Paroxysmal stage:</b> cough, usually coughing fits, characterized by inspiratory whoop (gasping) after repeated cough on the same breath, which commonly is followed by post-tussive emesis. Fever is absent or minimal. Older children may not have the whoop. Lasts 1–6 weeks, up to 10 weeks.  <b>Convalescent stage:</b> symptoms wane gradually over weeks to months. Lasts 2–3 weeks.	<b>PATIENT:</b> Exclude from school and child care until 5 days after initiation of appropriate antibiotic therapy. A 5-day course of azithromycin is usually treatment of choice and is also used for postexposure prophylaxis. Untreated individuals should be excluded until 21 days have elapsed from cough onset. If questions about pertussis treatment or postexposure prophylaxis, discuss with your local health department.  <b>CONTACTS:</b> PEP is recommended for all household and close contacts (including children in child care), people at high risk of severe pertussis (infants <12 months of age, people with certain pre-existing conditions), and those who will have contact with people who are high risk (including all people in child care settings) regardless of immunization status.  All close contacts should be monitored closely for respiratory tract symptoms for 21 days after last contact with the infected person. Close contacts with cough should have medical evaluation. Exclude on first indication of symptoms pending evaluation by healthcare provider.
<a href="#">Ringworm</a> (Body: Tinea Corporis, Scalp: Tinea Capitis, Groin: Tinea Cruris, Feet: Tinea Pedis)	1–3 weeks  Can be shorter, as reported cases have occurred at 3 days of age.	Skin-to-skin contact with lesions of infected persons or animals, shared objects (e.g., brushes, combs, hats, towels), shared surfaces (e.g., locker room floors), soil or fomites	Circular, red to brown, well-demarcated skin lesion(s). May have a scaly, vesicular, or pustular border with central clearing. Small confluent plaques or papules can occur. Itching is common.	<b>PATIENT:</b> Exclusion not indicated if child is receiving treatment. Child does not need to be sent home early from school but should begin treatment before returning. Cover lesions if able. (Scalp lesions do not need head coverings.)  <b>CONTACTS:</b> Exclusion is not indicated.

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

\*Please see last page for a summary of major changes.

<sup>†</sup>Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>



# Communicable Disease Reference Chart for School and Child Care Facility Personnel\*

DISEASE	INCUBATION PERIOD	TRANSMISSION	COMMON SYMPTOMS	RECOMMENDATIONS
<p><a href="#">Rubella</a><sup>†</sup> (German Measles)</p> <p><b>All suspected or confirmed cases of rubella are rapidly reportable to the local health department</b></p>	17 days (Range: 12–23 days)	<p>Direct or droplet contact from nasopharyngeal secretions.</p> <p>Most contagious when the rash is erupting, but communicable for 7 days before onset of rash and at least 7 days after onset of rash.</p>	Mild, maculopapular rash with swollen lymph nodes (often behind ear or at base of skull) and slight fever. Rash usually starts on face and spreads to the rest of the body within 24 hours. Rash typically lasts 3 days. Joint pain may occur, especially in older children and adults.	<p><b>PATIENT:</b> Exclude from school and child care for 7 days after onset of rash. Avoid exposure to women in early pregnancy. Check immunization records of all students. Discuss with your local health department.</p> <p><b>CONTACTS:</b> In outbreak setting, children without evidence of immunity should be immunized or excluded for 23 days after onset of rash of the last case in the outbreak. Pregnant contacts should be evaluated.</p>
<p><a href="#">Scabies</a></p>	<p>Persons without previous exposure: 4–6 weeks</p> <p>Previously infested and sensitized: 1–4 days</p>	<p>Prolonged close, personal contact. Infection from dogs/ animals is uncommon. Casual skin contact unlikely to result in transmission. Fomites are not likely to lead to classic scabies but can transmit crusted scabies because the parasite burden is much higher. (Crusted scabies is a severe, very contagious form that can occur in those with chronic conditions, developmental disabilities, or weakened immune systems.)</p>	Intensely itchy, red pimple-like rash around finger webs, wrists, elbows, knees, armpits, shoulder blades, waist, thighs, buttocks and/or genitalia. In children <2 years of age areas such as the scalp, face, neck, palms, and soles may be involved. Burrows are thin, gray or white, thread-like lines. Extensive scratching can result in secondary infection.	<p><b>PATIENT:</b> Exclude from school and child care until after the first course of appropriate treatment has been completed. Children identified during the school day should not be sent home early from school or child care because scabies has a low contagion within classrooms.</p> <p><b>CONTACTS:</b> Close contacts with prolonged skin-to-skin contact should receive treatment at the same time the infected person does. Bedding and clothing in contact with the skin of infected people should be laundered.</p>
<p><a href="#">Group A Streptococcal (GAS) Diseases</a> (Including Impetigo, Streptococcal pharyngitis “Strep throat”, and Scarlet Fever)</p>	<p>Impetigo: 7–10 days</p> <p>Strep throat/ Scarlet Fever: 2–5 days</p>	<p><b>Impetigo:</b> Direct contact with skin lesions or their discharge.</p> <p><b>Strep throat/Scarlet fever:</b> Respiratory droplets or contact with respiratory secretions. Fomite transmission and foodborne outbreaks are uncommon but can occur.</p>	<p><b>Impetigo:</b> Red, itchy sores that break open and leak clear fluid or pus, then develop a honey-colored crust. Commonly found on the arms, legs, mouth and nose.</p> <p><b>Strep throat:</b> Sudden onset of fever, sore throat, tonsil inflammation, palatal petechiae, strawberry tongue, and enlarged anterior cervical lymph nodes.</p> <p><b>Scarlet Fever:</b> A sandpaper-like rash that occurs with strep throat (or rarely with an infected wound).</p>	<p><b>PATIENT:</b> Impetigo: Exclude from school and child care until at least 12 hours after antibiotic treatment has started and all sores on exposed skin are covered. Close contact with other children should be avoided during this time.</p> <p>Strep throat: Exclude from school and child care until 12 hours after appropriate antibiotic therapy has been started AND afebrile without the use of fever-reducing medications (e.g. acetaminophen or ibuprofen).</p> <p><b>CONTACTS:</b> Impetigo: Exclusion not indicated.</p> <p>Strep throat: Symptomatic contacts should be medically evaluated and treated if positive. Routine lab testing or school/child care exclusion of asymptomatic household contacts is not indicated except during outbreaks or if the contact is at high risk of developing sequelae of infection. In general, chemoprophylaxis against Group A Strep is not recommended in school/child care settings.</p>

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

\*Please see last page for a summary of major changes.

<sup>†</sup>Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>

## Summary of Major Changes in the August 21, 2024 Update:

- Reviewed and updated content for all diseases using the Control of Communicable Diseases Manual, 21st Edition (2022) published by the American Public Health Association, the 2021-2024 Report of the Committee on Infectious Diseases, 32nd Edition (The Red Book) published by the American Academy of Pediatrics, and the relevant CDC website.
- Added link to Virginia Reportable Disease List in footer: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>.
- Updated information in the footer about consultation with the LHD, especially for outbreaks and reportable conditions, and added information on school-specific policies.
- Added link to CDC website for each illness, if available.
- Updated language from “immunocompromised” to “people with weakened immune systems”
- Chickenpox – updated to indicate that incubation period may be shorter in those with weakened immune systems, added more details about the rash, updated language regarding vaccination and exclusion of contacts
- COVID-19 – added language about variants, updated isolation and quarantine guidance
- Fifth Disease – added additional information about the onset of rash and infectious period
- Hepatitis A – updated exclusion criteria
- Hepatitis B – updated incubation period
- Influenza – clarified that recommendations only apply to seasonal influenza; added a note about novel influenza virus infections.
- Measles – updated incubation period with information about prodrome and rash onset
- Mpox – updated transmission information, exclusion criteria, and information on postexposure prophylaxis
- Mumps – updated contact exclusion criteria to indicate that students can be readmitted after receipt of a dose of MMR vaccine at the discretion of the health department.
- Pertussis – updated clinical description to include stages of illness; updated PEP recommendations to align with CDC
- Ringworm – Expanded from ringworm of the body to ringworm in general, which may include the scalp, groin, or feet. Updated exclusion guidance.
- Salmonellosis – updated incubation period
- Streptococcal Diseases – standardized language across the row to “strep throat” for “strep pharyngitis”

## Summary of Major Changes in the February 21, 2023 Update:

- Conjunctivitis (Pink eye) – Expanded from “acute bacterial” conjunctivitis to “all causes,” removed the requirement to be on antibiotics to return to school/child care, and added clearance by a healthcare provider to return while symptomatic.
- COVID-19 – Aligned transmission information with the Centers for Disease Control and Prevention (CDC). Removed information on close contacts.
- Fifth Disease (Parvovirus B19) – Added information about the characteristic rash.
- Hepatitis A – Clarified length of exclusion for patients who are food handlers and those who work in or attend child care.
- Mpox – Updated disease name from “monkeypox” to “mpox.”
- Pertussis – Aligned PEP recommendations for contacts with CDC recommendations.
- Rubella – Updated exclusion of unimmunized contacts from 21 days to 23 days.
- Streptococcal Diseases – Changed exclusion criteria from 24 hours of antibiotics to 12 hours of antibiotics for impetigo, and to 12 hours of antibiotics AND afebrile for strep pharyngitis.

NOTE: THESE RECOMMENDATIONS APPLY ONLY TO CHILDREN IN K-12 SCHOOLS OR CHILD CARE - A more complete discussion of these conditions and other communicable diseases may be found in *Control of Communicable Diseases Manual, 21<sup>st</sup> Edition* (2022) published by the American Public Health Association and the *2021-2024 Report of the Committee on Infectious Diseases, 32<sup>nd</sup> Edition (The Red Book)* published by the American Academy of Pediatrics. Additional information and consultation are also available through your local health department, especially for outbreaks and reportable conditions. School divisions may develop local policies for condition-specific recommendations.

\*Please see last page for a summary of major changes.

†Officially reportable in Virginia to the local health department. All disease outbreaks and unusual occurrences of disease are also reportable: <https://www.vdh.virginia.gov/content/uploads/sites/134/2023/03/VIRGINIA-REPORTABLE-DISEASE-LIST.pdf>